First Aid Test - Northeast Mine Rescue Contest - 2008

1.	Unless otherwise stated, all references to body when the body is in the			
	A. prone	C. lateral recumbent		
	B. supine	D. anatomical		
2.	The heart is located in	the	cavity.	
	A. thoracic	C. cranial		
	B. spinal	C. cranial D. abdominal		
3.	The primary purposes of the initial assessment is to:			
В. С.	<u> </u>	fe-threatening problems Il illnesses and injuries.		
4.	Respiration, circulatio	on, and bleeding are first	addressed during the:	
A.	focused history and p	hysical exam.		
	head-to-toe examination.			
	initial assessment.			
D.	taking of vital signs.			
5.	The carotid pulse can	be felt on the patient's:		
A.	neck			
В.	wrist			
	foot			
D.	temple			
6.	In the "ABCs" of emer	rgency care, the "C" star	nds for Compression.	
	True			
В.	False			

7.	7. A patient is short of breath and complains of pain in his left arm and jaw. You must care for this patient as through he may be experiencing:				
	A. congestive heart failure. B. a seizure	C. a heart attack. D. a stroke			
8. The primary responsibility of a First Responder providing care for a patient having a grand mal seizure is to place a bite stick in the patients mouth.					
	A. True B. False				
9.	9. The typical adult has about 12 pints of blood.				
A. B.	True False				
10. Bright red blood spurting from a wound indicatesbleeding.					
	A. venous C. capillary B. arterial D. ven	nule			
11.	11. Which of the following is a requirement for a nonemergency move?				
	A. The scene must be hazardB. The patient has a seriouslyC. You must reach another pD. The patient must be respo	y bleeding back wound. atient who is not breathing.			

	st commonly used de ack of the ambulance	0 -	patient, which is kept stretcher:			
	A. scoop B. flexible	C. wheeled D. basket (Stokes	s)			
13. Adequa A. True B. False	te blood flow to all co	ells of the body is ca	alled perfusion.			
14. The nor	14. The normal pulse rate for an adult at rest will fall between 60 and 100.					
A. True B. False						
15. Ineffect	15. Ineffective CPR will occur if:					
A. the breastbone is compressed.B. compression and release times are equal.C. the patient is on a hard surface.D. the patient's nostril are open during ventilation						
16. Except for a few special cases, impaled objects should:						
B. be re C. be re	oe removed moved if they are loo emoved if the impaler emoved to stop bleed	ment is shallow:				
	that involves all the burn.	layers of the skin is	commonly referred to			
_	al-thickness :hickness					

18. Which of the following chemicals should be brushed from the skin Rather than washed away with water?					
A. hydrochloric a B. dry lime	cid C. sodiur D. acetic	•			
19. An unknown chemical burn to the eye requires at leastminutes of flushing with running water.					
A. 5 B. 1	0 C. 15	D. 20			
20. A patient has a pain	20. A patient has a painful and deformed ankle injury. You should:				
A. straighten the foot and apply a pillow splint.B. tie his ankles together.C. Immobilize his foot in the position in which it was found, using a pillow splint.D. Elevate the entire lower extremity.					
21. When using a rigid splint on a suspected fractured femur, the longest splint should extend from theto past the foot.					
A. knee C. h B. buttocks D. v	-				
22. You can suspect possible brain injury if there is head trauma and the Pupils of the eyes are:					
A. responsive B. unequal	C. constricted D. pale red				
23. A patient with a head injury has clear fluids draining from the nose. You should:					
A. apply direct pressure with a dressing.B. pack the nostrils with dressing.C. pack the nostrils and ears with cotton.D. Not attempt to stop the flow.					

24. Your first concern for a patient who has and injury to the mouth tilt the head forward.	ı is to
A. True B. False	
25. If a patient has a head injury, you must suspect injures	ries.
A. spinal B. nasal	
C. eye D. ear	