Written First-Aid Examination 2010 Southern Regional Mine Rescue Contest New Iberia, LA

Name	Position
Team Name	

- 1. The thoracic cavity is also known as the chest cavity.
 - a. True
 - b. False
- 2. Which is <u>not</u> a guideline for ensuring a proper fit of a cervical collar?
 - a. Check to see it does not ride too far above or below the earlobes
 - b. Confirm the chin fits properly
 - c. Consider using a different size if it does not fit properly
 - d. Use the "two fingers" spacing method to avoid excessive tightness
- 3. If a patient is in the supine position, he is lying face up.
 - a. True
 - b. False
- 4. An emergency move should be considered when:
 - a. The patient may have spinal injuries
 - b. The patient is argumentative and may pose a hazard to rescuers
 - c. You are not able to gain access to other patients who need life-saving care
 - d. The patient has no visible blood loss.
- 5. The most important person during a log roll is the person at the patient's lower back/pelvic area.
 - a. True
 - b. False
- 6. Research by the American Heart Association suggests that the use of abdominal thrusts is still the most effective method for clearing the airway of a choking adult.
 - a. True
 - b. False
- 7. Which is <u>not</u> a step in assessing a patient's breathing?
 - a. Look for adequate tidal volume
 - b. Listen for air entering and leaving the nose and mouth
 - c. Observe the rise and fall of the lower abdominal area with each breath
 - d. Observe skin color

- 8. Which is not a recommended action for opening the airway?
 a. Reposition the head
 b. Jaw Thrust Maneuver
 c. Tilting the head and lifting the chin
 d. Head-Lift, Chin-Tilt Maneuver
 - 9. Which is <u>not</u> an example of an upper airway obstruction?
 - a. Obstruction by the tongue
 - b. Acute gastric distention
 - c. Mechanical obstruction
 - d. Obstruction by the epiglottis
- 10. You should not use the finger sweep on responsive patients or on unresponsive patients who have a gag reflex.
 - a. True
 - b. False
- 11. Which is <u>not</u> a major part of patient assessment?
 - a. Ongoing assessments
 - b. Scene size-up
 - c. Focused history and physical exam
 - d. Witness interviews
- 12. A slow, strong pulse may be indicative of what possible problem?
 - a. Overexertion
 - b. Heat emergency
 - c. Skull fracture
 - d. Internal bleeding (early stages)
- 13. A bulging neck vein is also known as a jugular vein distention.
 - a. True
 - b. False
- 14. The first component of the initial assessment is:
 - a. Assessing the patient's airway
 - b. Assessing the patient's mental status
 - c. Forming a general impression of the patient
 - d. Assessing the patient's circulation
- 15. The 1st rule for patient assessment is "Do No Further Harm".
 - a. True
 - b. False

- 16. Which is the most appropriate hand location for chest compressions on an adult?
 - a. At the top of the sternum
 - b. Two fingers above the breastbone line
 - c. Between the nipples over the lower half of the sternum
 - d. At the nipple line over the upper half of the sternum
- 17. A normal side effect of good chest compressions is the separation of the cartilage where the ribs meet the sternum.
 - a. True
 - b. False
- 18. When in doubt about a patient being hyperglycemic or hypoglycemic, follow the "sugar for everyone" rule.
 - a. True
 - b. False
- 19. Which is <u>not</u> a typical sign of ingested poisons?
 - a. Stains around the patient's mouth
 - b. Sweating
 - c. Abdominal pain
 - d. Excessive dry mouth
- 20. Which step does <u>not</u> apply when treating a patient with a generalized cold emergency?
 - a. Protect him from further heat loss
 - b. Monitor his vital signs
 - c. Do not allow the patient to walk
 - d. Provide warm liquids to drink
- 21. Which is <u>not</u> a category used to classify burns?
 - a. Partial thickness burn
 - b. Superficial burn
 - c. Full thickness burn
 - d. Secondary burn
- 22. Oddly enough, the loss of a tooth is considered an avulsion.
 - a. True
 - b. False
- 23. After the surface of an open wound is exposed, the next course of action should be to:
 - a. Control bleeding
 - b. Reassure the patient
 - c. Prevent further contamination
 - d. Remove superficial foreign matter with a sterile gauze pad

- 24. Which type of shock is seldom seen by emergency medical responders?
 - a. Hemorrhagic
 - b. Hypovolemic
 - c. Septic
 - d. Anaphylactic
- 25. In a sucking chest wound, the occlusive dressing should be taped on either three or four sides.
 - a. True
 - b. False
- 26. What is the order of care for skeletal injuries?
 - a. Spinal injuries, chest, skull, pelvis
 - b. Chest, spinal injuries, skull, pelvis
 - c. Spinal injuries, skull, pelvis, chest
 - d. Spinal injuries, skull, rib cage, pelvis
- 27. Which force is not a primary cause of musculoskeletal injury?
 - a. Direct force
 - b. Twisting force
 - c. Indirect force
 - d. Transitional force
- 28. What is the recommended order of steps to control bleeding?
 - a. Direct pressure, elevation, pressure points, tourniquet
 - b. Pressure points, direct pressure, elevation, tourniquet
 - c. Direct pressure, pressure points, elevation, tourniquet
 - d. Elevation, direct pressure, pressure points, tourniquet
- 29. When a patient suffers from angina pectoris, there is no actual damage to the heart muscle.
 - a. True
 - b. False
- 30. Which is <u>not</u> a primary tool for assessing a stroke victim when using the Cincinnati Prehospital Stroke Scale (CPSS)?
 - a. Facial droop
 - b. Arm drift
 - c. Memory loss
 - d. Abnormal speech