

# First-Aid Written Exam

Name \_\_\_\_\_

Team Name \_\_\_\_\_

Working Order \_\_\_\_\_

1. \_\_\_\_\_ mask would be the most important type of PPE to use when caring for a patient with tuberculosis.  
  - ☒ A. Hepa
  - B. Bag
  - C. Pocket
  
2. The abdominal cavity contains the liver and part of the \_\_\_\_\_ intestine.  
  - ☒ A. large
  - B. small
  - C. lower
  
3. When assessing circulation for a responsive adult patient you should assess the \_\_\_\_\_ pulse.  
  - A. femoral
  - B. carotid
  - ☒ C. radial
  
4. Skin that is \_\_\_\_\_ in color is called cyanotic  
  - A. red
  - ☒ B. bluish
  - C. pale
  
5. A patient who presents with normal vital signs and shows no indications of life-threatening problems may be described as \_\_\_\_\_.  
  - A. traumatic
  - B. normal
  - ☒ C. stable

## First-Aid Written Exam

6. Your patient has been in respiratory distress for approximately 30 minutes, your assessment reveals pale skin and cyanosis of the lips, these are signs of \_\_\_\_\_
- A. bronchitis
  - B. asthma
  - ☒ C. hypoxia
7. Protect the patient from injury and place him or her in the \_\_\_\_\_ following the seizure is an example of appropriate care for a seizure patient.
- ☒ A. Recovery position
  - B. Prone position
  - C. Supine position
8. A wound where the top layers of skin have been scraped off, commonly seen in falls, can best be described as an \_\_\_\_\_
- A. contusion
  - ☒ B. abrasion
  - C. laceration
9. When the \_\_\_\_\_ is absent is a situation where it would be appropriate to place an angulated extremity back into the anatomical position.
- ☒ A. Distal pulse
  - B. Radial pulse
  - C. Carotid pulse
10. You are caring for a patient who has one leg that is shortened with the foot rotated to one side, these are likely signs of a possible \_\_\_\_\_
- A. Dislocated femur
  - B. Dislocated knee
  - ☒ C. Dislocated hip

## First-Aid Written Exam

11. The most appropriate care for an open abdominal injury is to cover the wound with a \_\_\_\_\_, sterile dressing
- A. clean
  - ☒ B. moist
  - C. dry
12. A ratio of \_\_\_\_ compressions per minute and a depth of at least 2 inches are the rate and depth for chest compressions on an adult.
- A. 80 to 100
  - ☒ B. 100 to 120
  - C. 30 to 2
13. When more rescuers arrive on scene you should assign tasks to other rescuers and rotate compressors every \_\_\_\_ or more frequently if needed to avoid fatigue.
- ☒ A. 2-minutes
  - B. 5-minutes
  - C. 3-minutes
14. \_\_\_\_\_ is an important tool for identifying whether opioids may be involved in a life-threatening emergency.
- A. Patient assessment
  - B. Initial assessment
  - ☒ C. Scene assessment
15. \_\_\_\_\_ is a life-threatening condition that occurs when the circulatory system can't maintain adequate blood flow.
- A. stroke
  - ☒ B. shock
  - C. asthma